International Journal of Trend in Scientific Research and Development (IJTSRD)
Volume 10 Issue 5, Sep-Oct 2026 Available Online: www.ijtsrd.com e-ISSN: 2456 – 6470

Erectile Dysfunction: A Review with a Homoeopathic Perspective

Dr. Kishan Adesara[1], Dr. Riddhish Tanna[2], Dr. Nishita Doshi[3]

[1]PG Scholar, [2]MD Homoeopath (Professor), [3]MD Homoeopath (Associate Professor),
[1,2,3]Shree B.G. Garaiya Homoeopathic Medical College & Hospital, Saurashtra University, Rajkot, Gujarat, India

ABSTRACT

Erectile dysfunction (ED) is a common male sexual health condition characterized by the persistent or recurrent inability to attain or maintain a penile erection sufficient for satisfactory sexual performance. It may be associated with vascular, metabolic, hormonal, neurological, psychological, and lifestyle-related factors. Erectile dysfunction can adversely affect self-esteem, intimate relationships, psychological well-being, and overall quality of life. Despite its clinical importance, embarrassment, social stigma, and reluctance to discuss sexual concerns may delay diagnosis and treatment. This review summarizes the definition, aetiology, pathophysiology, clinical manifestations, diagnostic evaluation, and contemporary management of erectile dysfunction. It also discusses homoeopathic concepts of individualization, totality of symptoms, susceptibility, and miasmatic interpretation as a theoretical approach to case-taking. The role of individualized assessment and appropriate medical evaluation is emphasized. Scientific evidence supporting homoeopathy specifically for erectile dysfunction remains insufficient; therefore, it should not replace established diagnostic evaluation or evidence-based treatment.

KEYWORDS: *Erectile dysfunction; male sexual health; homoeopathy; individualization; totality of symptoms; susceptibility; miasmatic interpretation; quality of life.*

How to cite this paper: Dr. Kishan Adesara | Dr. Riddhish Tanna | Dr. Nishita Doshi "Erectile Dysfunction: A Review with a Homoeopathic Perspective" Published in International Journal of Trend in Scientific Research and Development (ijtsrd), ISSN: 2456-6470, Volume-10 | Issue-5, October 2026, pp.1791-1794, URL: www.ijtsrd.com/papers/ijtsrd142298.pdf

IJTSRD142298

Copyright © 2026 by author (s) and International Journal of Trend in Scientific Research and Development Journal. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0) (http://creativecommons.org/licenses/by/4.0)

INTRODUCTION

Erectile dysfunction is defined as the consistent or recurrent inability to attain and/or maintain a penile erection sufficient for satisfactory sexual performance. It is an important clinical concern because it may affect physical health, psychological well-being, intimate relationships, and quality of life. [1,2]

Penile erection is a complex physiological process involving psychological stimulation, neurological pathways, vascular mechanisms, hormonal influences, and relaxation of cavernosal smooth muscle. Disturbance in any of these components may contribute to erectile dysfunction.

ED may occur in association with diabetes mellitus, hypertension, obesity, cardiovascular disease, hormonal abnormalities, neurological disorders, certain medications, anxiety, depression, and relationship difficulties. It is not an inevitable consequence of ageing and may occur in younger as well as older men. [2,3]

Although effective treatment options are available, some individuals hesitate to seek medical advice because of embarrassment, cultural expectations, fear of judgement, or concerns about sexual performance. Early assessment is important because ED may also indicate underlying cardiovascular or metabolic disease. [1,2]

This review summarizes current medical understanding of erectile dysfunction and discusses homoeopathic concepts relevant to individualized case-taking and theoretical interpretation.

ETIOLOGY AND PATHOPHYSIOLOGY

Erectile dysfunction is a multifactorial condition. Its causes may be broadly classified as follows:

1. Vascular factors: Penile erection depends on adequate arterial blood flow and effectiverestriction of venous outflow. Endothelial dysfunction, atherosclerosis, hypertension, and cardiovascular disease may impair the erectile response.

2. Metabolic factors: Diabetes mellitus, obesity, insulin resistance, and dyslipidaemia may contribute to vascular and neurological changes associated with ED.
3. Hormonal factors: Low testosterone and certain endocrine disorders may affect sexual desire and erectile function. Hormonal abnormalities should be evaluated when clinically indicated.
4. Neurological factors: Conditions affecting the central or peripheral nervous system, spinal cord injuries, and complications of pelvic surgery may interfere with the neural pathways responsible for erection.
5. Psychological factors: Performance anxiety, stress, depression, fear offailure, and relationship difficulties may contribute to erectile problems. Psychological and organic factors can coexist.
6. Medication and substance-related factors: Certain prescribed medicines, tobacco use, excessivealcohol consumption, and recreational substances may contribute to sexual dysfunction. Medication-related concerns should be reviewed by a qualified clinician.
7. Lifestyle-related factors: Physical inactivity, inadequate sleep, obesity, and unhealthy dietary patterns may contribute to ED through their effects on metabolic and cardiovascular health. [1–3]

CLINICAL CLASSIFICATION

Erectile dysfunction may be described according to its predominant cause:

- Organic ED: Associated with vascular, neurological, hormonal, anatomical, or other physical factors.
- Psychogenic ED: Predominantly associated with psychological or situational factors.
- Mixed ED: Involves a combination of physical and psychological contributors.

These categories are not always mutually exclusive. A comprehensive assessment is necessary to identify the factors relevant to an individual patient.

CLINICAL PRESENTATION

Clinical manifestations may include:

- Difficulty attaining an erection.
- Difficulty maintaining an erection during sexual activity.
- Reduced erectile firmness.
- Inconsistent erectile responses in different situations.
- Reduced sexual confidence and increased performance anxiety.
- Emotional distress, relationship difficulties, or reduced quality of life.

The presence or absence of morning erections, erections during masturbation, and situational variation may help inform clinical assessment, but these features alone cannot establish the cause of ED. [1,2]

DIAGNOSTIC EVALUATION

Evaluation should include a detailed medical, sexual, and psychosocial history, a focused physical examination, and appropriate investigations. [1–3]

1. Medical and sexual history: Assessment may include the onset, duration, frequency, severity, and circumstances of erectile difficulty; sexual desire; morning erections; previous illnesses; medications; substance use; and relevant surgical history.
2. Psychological and relationship assessment: Stress, anxiety, depressive symptoms, performance concerns, and relationship factors should be explored sensitively and without judgement.
3. Physical examination: Blood pressure, body mass index, cardiovascular risk factors, and relevant genital and endocrine findings may be assessed according to the clinical presentation.
4. Laboratory investigations: Testing may include fasting glucose or HbA1c, lipid profile, and morning total testosterone. Additional hormonal or other investigations may be indicated by the history and examination.
5. Standardized assessment tools: Validated instruments such as the International Index of Erectile Function (IIEF) and the Sexual Health Inventory for Men (SHIM) may help assess severity and monitor changes over time. [1,3]

MANAGEMENT

Management should be individualized according to the underlying cause, medical history, patient preferences, and treatment safety.

Lifestyle modification: Regular physical activity, weight management where appropriate, smoking cessation, moderation of alcohol consumption, adequate sleep, and a balanced diet may improve general health and support erectile function.

Medical treatment: Phosphodiesterase type 5 inhibitors, including sildenafil and tadalafil, are established treatment options for appropriate

patients. They require clinical assessment and may be contraindicated with nitrate medications because of the risk of a dangerous fall in blood pressure.

Management of underlying conditions: Diabetes, hypertension, dyslipidaemia, cardiovascular disease, and hormonal disorders should be evaluated and managed appropriately.

Psychological and psychosexual support: Counselling or psychological therapy may be beneficial when anxiety, depression, performance concerns, or relationship difficulties contribute to ED.

Other treatment options: Vacuum erection devices, selected injectable therapies, and penile prostheses may be considered in suitable cases when clinically indicated. Treatment decisions should be made through informed discussion with a qualified healthcare professional. [1,3]

HOMOEOPATHIC PERSPECTIVE

UNDERSTANDING THE INDIVIDUAL PATIENT

Classical homoeopathic philosophy emphasizes understanding the individual patient rather than selecting a remedy solely on the basis of a diagnostic label. In the context of erectile dysfunction, this approach would involve documenting the patient's physical complaints, emotional experiences, general characteristics, personal history, and relevant circumstances.

However, this philosophical approach should be distinguished from established clinical evidence. There is insufficient reliable evidence to conclude that homoeopathic remedies effectively treat erectile dysfunction.

INDIVIDUALISATION

Individualization in homoeopathic practice refers to considering the characteristic features of the individual case. A case record may include:

- Onset and progression of erectile difficulty.
- Ability to attain and maintain an erection.
- Sexual desire and relevant sexual history.
- Morning and situational erections.
- Emotional state, anxiety, and stress.
- Sleep, appetite, thirst, and general physical characteristics.
- Relevant medical conditions, medications, and investigations.
- The patient's concerns, expectations, and treatment priorities.

These details can support a comprehensive clinical history. They must not substitute for medical evaluation or delay appropriate treatment.

TOTALITY OF SYMPTOMS

In homoeopathic literature, the totality of symptoms refers to the complete individual symptom picture considered during remedy selection. In ED case-taking, this may encompass the presenting complaint, associated physical symptoms, psychological experiences, modalities, medical history, and general characteristics.

The clinical significance of each feature should be assessed carefully, and symptoms should not be interpreted as proof of a particular underlying pathology without appropriate investigation.

SUSCEPTIBILITY

Susceptibility is a concept in classical homoeopathic literature used to describe an individual's perceived responsiveness to influences. Within a theoretical case analysis, practitioners may document general health, the duration and intensity of complaints, associated conditions, previous treatment, and individual responses.

This concept should be understood as part of homoeopathic philosophy rather than a validated biomedical measurement of erectile function.

MIASMATIC INTERPRETATION

Within classical homoeopathic philosophy, chronic diseases have traditionally been interpreted through the concepts of psora, sycosis, and syphilis, described in the works of Samuel Hahnemann, including The Chronic Diseases and Organon of Medicine.

In a theoretical homoeopathic analysis of erectile dysfunction, the following interpretations may be discussed:

Psoric interpretation: Functional complaints, anxiety, insecurity, and perceived sensitivity may be explored within a traditional psoric framework when supported by the broader individual symptom picture.

Sycotic interpretation: Features interpreted by classical homoeopathic practitioners as excessive, persistent, or associated with a particular pattern of chronic complaints may be discussed within a sycotic framework.

Syphilitic interpretation: Destructive or degenerative features, where relevant to the patient's broader clinical history, may be considered within traditional syphilitic interpretation.

Mixed miasmatic interpretation: A practitioner may describe a mixed miasmatic picture when the recorded

characteristics are interpreted as corresponding to more than one traditional category.

These interpretations are theoretical constructs within homoeopathic philosophy. They are not scientifically validated classifications of erectile dysfunction, and no single miasmatic category can be assigned to all patients with ED. Erectile dysfunction should be assessed according to established medical criteria, and its management should not be based on miasmatic interpretation alone.

CONCLUSION

Erectile dysfunction is a multifactorial condition that may arise from vascular, metabolic, hormonal, neurological, psychological, and lifestyle-related factors. Its effects may extend beyond sexual performance to psychological well-being, self-confidence, intimate relationships, and overall quality of life. A comprehensive history, physical examination, appropriate investigations, and individualized evidence-based management are essential for identifying contributing factors and selecting suitable treatment.

Homoeopathic concepts such as individualization, totality of symptoms, susceptibility, and miasmatic interpretation provide a traditional philosophical framework for case-taking. However, the clinical effectiveness of homoeopathy for erectile dysfunction has not been established by sufficient reliable evidence. Further well-designed clinical research is required to determine whether individualized homoeopathic treatment offers any benefit beyond placebo. Conventional medical evaluation and evidence-based treatment should not be delayed or replaced.

REFERENCES

[1] Burnett AL, Nehra A, Breau RH, Culkin DJ, Faraday MM, Hakim LS, et al. Erectile dysfunction: AUA guideline. J Urol. 2018; 200(3): 633-641. doi:10.1016/j.juro.2018.05.004.

[2] National Institute of Diabetes and Digestive and Kidney Diseases. Erectile dysfunction (ED) [Internet]. Bethesda (MD): National Institutes of Health. Available from: https://www.niddk.nih.gov/health-information/urologic-diseases/erectile-dysfunction

[3] European Association of Urology. EAU guidelines on sexual and reproductive health: management of erectile dysfunction [Internet]. Arnhem: European Association of Urology. Available from: https://uroweb.org/guidelines/sexual-and-reproductive- health/chapter/management-of-erectile-dysfunction

[4] Hahnemann S. Organon of Medicine. 6th ed. Boericke W, translator. Bibliographic details should be completed according to the edition consulted.

[5] Hahnemann S. The Chronic Diseases: Their Peculiar Nature and Their Homoeopathic Cure. Bibliographic details should be completed according to the edition consulted.