Enrichment of Omega-3 Fatty Acids using Urea Complexation Method to Enhance the Nutritive Value of Stingray Fish (Dasyatis Sephen F.) Liver Oil

Duglas Sathees¹, Vidanarachchi J. K², Himali S. M. C²

¹Department of Animal Science, Faculty of Agriculture, University of Jaffna, Sri Lanka ²Department of Animal Science, Faculty of Agriculture, University of Peradeniya, Sri Lanka

How to cite this paper: Duglas Sathees | Vidanarachchi J. K | Himali S. M. C "Enrichment of Omega-3 Fatty Acids using Urea Complexation Method to Enhance the Nutritive Value of Stingray Fish (Dasyatis Sephen F.) Liver Oil" Published

International in Journal of Trend in Scientific Research Development and (ijtsrd), ISSN: 2456-6470, Volume-3 | Issue-5, August 2019, pp.24-29,



https://doi.org/10.31142/ijtsrd25181

Copyright © 2019 by author(s) and International Journal of Trend in Scientific Research and Development Journal. This is an Open Access article distributed

under the terms of the Creative **Commons Attribution**

License



(CC (http://creativecommons.org/licenses/by (4.0)

This paper describes the study performed on Ray fish liver oils from fish by-products discarded from a dry fish processing industry where usually discarded.

Even though refined fish oil has a numerous health benefits, only certain people afford to consume it as fish oil is an expensive item. Presently, the fish production is becoming more demanding as there is a sizable and growing world market demand for high quality fish oil. Apart from its various uses as consumable oil, it is also appreciable in both pharmaceuticals and food industries. Past literature has shown that little work has been done in terms of Ray fish liver oil and its fish by-products. Therefore, Ray fish liver oil may have applications in food, feed manufacturing and other industries via improve nutritive value of crude ray fish live oil.

2. Materials and Methods

2.1 Study Location

This study was conducted in the Fisheries Harbors of Gurunagar and Delft Islands, Jaffna, Northern Provenience in Sri Lanka. Analysis was conducted in the Nutrition Laboratory, Meat Science Laboratory, Department of Animal Science, Food Science Laboratory, Department of Food Science, Faculty of Agriculture, and University of Peradeniya and in the Beuro Veritas, Colombo, Sri Lanka.

ABSTRACT

Lipid fraction extracted from tissues of oily fish and fishery by-products are one of the best source of omega-3 (ω -3) polyunsaturated fatty acids (PUFAs), mainly eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Method of oil extraction from the Dasvatis sephen liver is simple and cheap. Therefore, the present study was conducted to extract and enrich the Omega-3 fatty acids from Dasyatis sephen liver which is discarded during dry fish production. Bligh and Dyer's method was used to extract oil. Fatty acids profiles were determined by Gas-Liquid Chromatography method (GLC). Average liver lipid recovery was 69.54 % (w/w). Crude liver oil fatty acid profile of DHA and EPA was 0.5% and 0.6%, respectively. Urea complexation was done to enrich the extracted Omega-3 fatty acids. Physio-chemical properties such as moisture content, color, specific gravity, peroxide value, and fatty acid compositions were obtained under the tolerable standard. The level of DHA:22:6n-3 and EPA: 20:5n-3 in the enriched Dasyatis sephen oil were 9.7 % and 8.7 %, respectively. It could be concluded that the converting of Ray fish by-product into enriched oil is an opportunity of adding value to the fish by-product and could be suitable for applications in pharmaceutical and nutraceutical industries.

KEYWORDS: Fatty acid, Fish Oil, Omega-3, Ray Fish, Urea Complexation

1. INTRODUCTION

Fish oil is the lipid fraction extracted from tissues of oily fish and fishery byproducts and it is the best source of omega-3 (ω -3) polyunsaturated fatty acids (PUFAs), especially eicosapentaenoic acid (EPA) and docosahexaenoic acid

(DHA).

2.2 Sample Collection and Preparation

Samples of Three Stingray fish Dasyatis sephen (Forsskal, 1175) were collected from each Fisheries Harbors and Dry Fish Cottages of Jaffna. They were transported in sealed freezer containers to laboratory at (-8 °C) and stored at (-20 °C) in the deep freezer for analyses. Each were taken out to thaw at room temperature to measure its lengths and weights as whole. Samples were eviscerated and its livers, offal (gut & tail) and carcass were stored at (-20 °C) in sealed in poly bags separately for further analysis.



Plate 2.1: Sting Ray (Dasyatis sephen) (Dorsal View)

2.2.2 Extraction of Ray Fish Liver Crude Oil

A modified Bligh and Dyer method (1959) was used to determine the crude lipids in the liver samples of ray fish. Extraction of liver oil was carried out by solvent extraction method. 50 g of ray fish liver samples were homogenized for two minutes using rotary homogenizer Bio-Gen PRO200 with 100 mL of methanol and 50 mL of chloroform. Then, another 50-mL of chloroform was added and homogenized using the homogenizer for another 30 Seconds. The mixture was diluted with 50 mL of distilled water.

Then mixtures were filtered through a Buncher funnel lined with Whatman No.42 filter paper under vacuum suction. The residue was rinsed using 20 mL of chloroform and the filter was transferred to a separator funnel. The aqueous layer was removed and the chloroform layer containing ray fish liver oil (bottom layer) was separated. It was filtered using a Buncher funnel lined with Whatman No.1 filter paper under vacuum suction. The residue was again rinsed with 20 mL of chloroform. The filtrate was passed through 3 g of anhydrous sodium sulfate to remove moisture.

Chloroform layer was transferred into a dry pre-weighed round-bottom flask. The chloroform was evaporated using a rotary vacuum evaporator at 40 °C. Extracted crude liver oil was weighed and treated with 0.02 BHT. Finally the prepared crude liver oil was stored at (-20 °C).



Plate: 2.2 Fish oil Colour

2.2.3 Crude Oil Recovery Percentage

Below given equation was used to calculate the lipid content in the sample.

Equation: Percentage of lipid in the Ray fish liver fish

% Lipid =
$$\frac{W_L}{W_S}$$
 X 100

W_S= Weight of the sample (g)

W_L= Weight of Lipid extracted after evaporation (g)

2.2.4 Determination of fatty acid composition in Crude oil

Fatty acids profiles of selected oils were determined according to Buchgraber *et al.* (2000) method by Gas-Liquid Chromatography (GLC).

2.2.5 Preparation of free fatty acid for urea complexation

25 g of Ray fish liver oil was measured into a 250 ml perdried conical flask. A mixture of 95% ethanol of 66 ml, distilled water of 11ml and KOH of 5.75g in pellet weight were added to the oil. The solution was refluxed at (-8.5 +/-2.0 degree celsius under nitrogen blanket for 1 day. Then the solution was diluted with 50 ml of diluted water and mixed well. 100 mL of n-hexane was added and mixed well.

The Mixture was transferred to a separator funnel and aqueous layer containing saponified matter was separated. The n-hexane layer containing unsaponified matter was discarded. The separated aqueous layer was washed again using 100 mL of n-hexane. The n-hexane layer on top was separated by separator funnel and was discarded. The aqueous layer in bottom was acidified using 3 N HCl solution until pH reached 1. Then the solution was mixed with 50 mL of hexane and transferred to the separator funnel. The n-hexane layer containing fatty acid was separated and treated with 3 g of anhydrous sodium sulfate. Finally n-hexane was evaporated at 40 °C using a rotor evaporator. Liberated free fatty acid was weighed and stored at (-20 °C) after filling into 20 mL labeled glass vials.

2.2.6 Preparation of Omega- 3 concentrates using urea complexation

100 ml of 95% aqueous ethanol was added. The required urea amount was calculated according to the free fatty acid weight, urea to free fatty acid ratio was maintained at 6:1. The system was heated to 65°C and continuous stirring was practiced until the solution becomes crystal clear amber color solution. Then the system was rapidly transferred to a refrigerator where the temperature was adjusted to (- 9) °C and kept for 18 hours in the freezer FRIMED S.R.L AF70V/2 for crystallization.

> After 18 hours, the mixture was allowed to reached room temperature 27°C. Then the fraction was separated by filtering using a Bunchner Funnel lined with Whatman No.1 filter paper. The filter is known as Non-Urea Complexing Fraction. The residue is the urea complexing fraction.

The Fraction containing Non-Urea Complexation was diluted with an equal volume of distilled water and acidified to pH 4 – 5 using freshly added 6 N HCl solutions. An equal amount of n-hexane was added and the mixture was stirred thoroughly for 1 hour using a magnetic stirrer. Then, the solution was transferred to a separator funnel to separate nhexane layer. The n-hexane layer was treated with 3 g of anhydrous sodium sulfate to remove moisture. Vacuum facilitated filtration was practiced to remove hydrated sodium sulfate using a Buchner Funnel lined with Whatman No.1 paper. The n-hexane was evaporated using a rotary evaporated. The liberated Omega–3 fatty acids were treated with 0.02g BTH and stored in (-20°C) until analysis.

2.2.7 Preparation of fatty acid methyl ester (FAME)

Sodium methoxide (CH₃NaO) (0.5M): 0.074 g sodium methoxide was weighed accurately and dissolved in 2 mL of methanol. Internal standard (Methylheptadecanoate) (1mg/mL): 100 mg of methylheptadecanoate acid was dissolved in 100 ml of hexane.

Sample (100 mg) was weighed in to a 15 mL screw capped methylation tube and 1 mg/mL of internal standard

(Methylheptadecanoate), 2mL of 0.5M sodium methoxide and 300 UL of dichloromethane were added. Subsequently, mixture was kept in a heat block at 50°C for 30min. It was allowed to cool to room temperature and 5mL of distilled water added drop by drop and 100 UL of glacial acetic was added. Then the tube with contents will be centrifuged at 1500 rpm for 10 minutes at 5°C and top hexane layer will be separated and added in to a 2ml GC Vial. Then against 500 UL of hexane will be added and centrifuged as mentioned above and top hexane layer will be separated and added to the same GC vial. Vials will be sealed with Para film and frozen immediately at (-20°C) until analysis by GC.

2.2.8 Gas Liquid chromatography analysis of sample

The prepared FAME was analyzed by injecting 1 µL into GLC (Shimadzu, 14-B, Japan), equipped with a Flame Ionization Detector (FID) and a fused silica capillary column (100 m, 0.25 mm id and 0.20 µm film thickness) and attached with Chromotopac data processor (Model-CR6A, Shimadzu, Japan). Split ratio was 100:1. Injector and detector temperatures were maintained at 260°C. Helium was used as carrier gas at flow rate of 20 mL/sec. The initial column oven temperature was maintained at 140°C for 5 min and increased to 220°C at the rate of 4°C/min, then maintained at that temperature for 10 min. Fatty acids were identified by comparison of their retention time with authentic standards (SUPELCO 37 Component FAME Mix, Sigma Aldrich) (Kuksis *et al.*, 1967). The amount of each fatty acids in the sample.

3. Results and Discussion

3.1 Hepatosomatic Index (HSI) of Ray fish liver
The average body weights, average lengths and Liver weight of the String fish were found to be 2301.05g, 2594.17g and 2847.84g where lengths were 137.67cm, 153.49cm and 175.74cm. The liver weight were 253.69g, 289.52g and 313.98 g respectively. HSIs were calculated to be 9.05%. Moreover the average lengths of stingray were measured to be 137.61 and 148.67 cm, 171.32 cm respectively. The HSI value of String ray fish is comparatively greater than the findings of Özyılmaz and Öksüz, 2015 which is 8.25%. This increment may be due to the increased liver weight of String ray fish and the maturity level of fish.

Table3.1: The average length (cm), Total weight (g), Liver weight (g) and Hepatosomatic index (HSI) of the Bay fish

Fish species	Length (cm)	Total weight (g)	Liver weight (g)	HSI (%)
Stingray-1	137.67	2301.05	253.69	9.07
Stingray-2	153.49	2594.17	289.52	8.96
Stingray-3	175.74	2847.84	311.92	9.13

Mean±SD values were presented (n = 3)

3.2 Stingray Fish Liver Crude Oil Recovery Percentage The average liver lipid levels of the String ray fish was 69.54 %. All of the ray fish had a large amount of lipids in their liver. However, the levels of liver lipid not varied greatly (Özyılmaz and Öksüz, 2015).

3.3 Fatty acid Profile of Ray fish crude liver oil

The predominant fatty acid in the stingray MUFA was C16 (palmitic acid), which had a value of 35.0%. Clearly, the average level of C16 in the ray fish liver oil was significantly

higher than the level of the other ray fish liver oils. The total levels of PUFA in all the fish liver oils seems similar.

The total SFA percentages of String ray fish in this study were 44.2%. The total SFA percentages of ray fish liver oils from String ray that were previously reported were 34.97%, respectively (Navarro-Garcia *et al.*, 2010). The percentages of total SFA in ray fish liver oil was not in agreement with the finding from the current study.

Table3.2: Fatty acid composition of stingray fish liver

011				
Carbon Chain	Stingray			
C12:0	4.1			
C13:0	0.7			
C14:00	0.3			
C16:0	35.0			
C17:0	0.4			
C18:0	2.5			
C20:0	1.2			
ΣSFA	44.2			
C16:1	10.2			
C18:1	1.3			
ΣMUFA	11.5			
C18:2	0.3			
C20:4	1.4			
Σn6	1.7			
C20:5n3	0.5			
C22:6n3	0.6			
Σn 3	1.1			
ΣΡυΓΑ	2.8			
n3/n6	0.64			
O DHA/EPA	0.83			

3.4 Fatty acid Profile of Enriched Ray fish liver oil

The predominant fatty acid in the enriched sting ray MUFA was C16 (palmitic acid), which had a value of 15.2%. Clearly, the average level of C16 in the ray fish liver oil were significantly lower than the level of the other ray fish liver oils. The total levels of PUFA in all the enriched ray fish liver oils increased significantly.

The total SFA percentages of enriched String ray fish in this study were 28.8%. The total SFA percentages of two ray fish liver oils from String ray that were previously reported was 34.97%, respectively (Navarro-Garcia *et al.*, 2010). The percentages of total SFA in fish were not in agreement with the finding from the current study.

In this study, the level of Σ MUFA 11.5% in the liver oil of the stingray was lower than the levels found in string ray.(Ould El Kebir *et al.*, 2005). Surprisingly, in contrast to containing high levels of C16, the liver oil of the ray fish has high amounts of EPA, and DHA, which are not generally dominant in marine fish.

In this study, the levels of DHA and EPA in ray fish liver oil differed greatly from the enriched ray fish oil. The levels of DHA and EPA in enriched ray liver oil were 9.7 % and 8.7 %, which is heigher than the levels of DHA and EPA for male *Dasyatis marmorata* that Ould El Kebir *et al.* (2007) reported.

Table 3.3: Fatty acid composition of Enriched stingray

ray fish liver oil			
Carbon Chain	Stingray		
C12:0	1.1		
C13:0	0.0		
C14:00	4.0		
C16:0	15.2		
C17:0	0.7		
C18:0	4.2		
C20:0	3.6		
ΣSFA	28.8		
C16:1	18.5		
C18:1	6.3		
ΣMUFA	24.8		
C18:2	1.3		
C20:4	2.3		
Σn6	3.6		
C20:5n3	8.7		
C22:6n3	9.7		
Σn3	19.8		
ΣΡυγΑ	23.4		
n3/n6	5.5		
DHA/EPA	0.89		

4. Conclusions

Ray fish by-product (liver) was obtained during dry fish processing of ray fish at Jaffna, Northern province, Sri Lanka. The liver oil was obtained through the Bligh and Dyer method gave the total recovery percentage of the ray fish liver oil was 69.54 %.. The crude enriched via urea complexation method enhance the nutritive value of ray fish liver oil by enriched ray fish liver oil with high in omega-3 in Scie with added antioxidants: Effects of temperature and content of 9.7 % and 8.7 %. Enriched ray fish liver oil from dry fish industry by-product could be suitable for applications as nutraceutical in the food industry.

5. References

- [1] Ackman RG. Fatty Acid Composition of Fish Oils, Nutritional Evaluation of Long Chain Fatty Acids in Fish Oil. Edited by SM. Barlow and ME. Stansby, Academic Press, London, 1982, 25-88.
- [2] Ackman RG. Fish in Marine Biogenic Lipids, Fats and Oils. CRC Press, Inc., Boca Raton 1989; 2:145-178.
- [3] Ackman RG. Nutritional Composition of Fats in Sea foods. Prog Food Nutr Sci 1989; 13:161-241.
- Ackman, R.G. 2005. Marin lipids and omega-3 fatty [4] acids. In: Akon, C.C. (ed.), Handbook of Functional Lipids. Taylor and Francis Group, New York, USA, 311-24.
- [5] Aidos, I., J.B. Luten, R.M. Boom and A.V. Padt. 2001. Upgrading of Maatjes herring byproducts: Production of crude fish oil. Journal of Agricultural Food Chemistry, 49,3697-3704
- [6] Aidos, I., S. Masbernat-Martinez, J.B. Luten, R.M. Boom and A.V. Padt. 2002. Composition and stability of herring oil recovered from sorted by-products as compared to oil from mixed by-products. Journal of Agricultural Food Chemistry, 50, 2818-2824.
- [7] Aidos, .I, Van Der P.A., Luten, J.B, Boom, R.M. 2002. Seasonal Changes in Crude and Lipid Composition of Herring Fillets, Byproducts, and Respective Produced Oils. Journal of Agricultural and Food Chemistry 16: 4589-4599.

- [8] Alders, L. 1955. Liquid-liquid extraction. Elsevier, New York, 26
- Bharadwaj. 2011. Synthesis of DHA rich PUFA from cod [9] liver fish oil. http://www.scribd.com/doc/70347867/Synthesis-of-DHA-Rich-PUFA-From-CodLiver-Fish-Oil.
- [10] Bimbo, A.P., 1998. Guidelines for characterizing foodgrade fish oils. Inform. Int. News Fats Oils Relat. Mater., 9(5): 473-483.
- Bimbo, A.P. (2007). Processing of marine oils. In H. [11] Breivik (Ed.), Long-chain omega-3 specialty oils (pp. 77–109). Oily Press Lipid Library. Bridgwater, UK: P.J. **Barnes & Associates**
- Bligh, E.G. and W.J. Dyer. 1959. A rapid method for total [12] lipid extraction and purification. Canadian Journal Biochemistry and Physiology, 37, 911-917
- [13] Boran, G., H. Karacam and M. Boran. 2006. Changes in the quality of fish oils due to storage temperature and time. Food Chemistry, 98, 693-698.
- Boxshall, G.A.; Mees, J.; Costello, M.J.; Hernandez, F.; [14] Gofas, S., et al. 2014. World Register of Marine Species.
- British Nutrition Foundation. 1992. Unsaturated fatty [15] acids. Nutritional and Physiological significance. Report of British Nutrition Foundation. Chapman & Hall, London 156-157
- Bruinsma, K.A. and D.L. Taren. 2000. Dieting, essential [16] fatty acid intake, and depression, 58(4), 98-108.
- [17] onal J Chingmin, Ph.D., E. Tsai, T. Joseph, Wooten, Ph.D and D.A. Otto. 1989. Stability of fish oil in a purified diet arch an light. Nutrition Research, 9(6), 673-678.
- evelo [18] Christian A. Drevon.2010. Omega-3 fatty acidsmetabolism and mechanisms of action of essential fatty ISSN: 2456-647 acids. ISBN-978-82-303-1590-3.
 - Christie, W.W. 2003. Lipid extraction, storage and [19] sample handling. In: Lipid analysisisolation, separation, ... identification and structural analysis of lipids (3rd edition), Christie (ed). Oily Press, England, 97-10.
 - [20] Chol Su Pak2005. Stability And Quality Of Fish Oil. During Typical Domestic Application. Wonsan University of Fisheries
 - .Debasish P, Dipankar B, Tarun KP, Amarendra [21] P,Amitabha G. Liver lipids and fatty acids of the sting ray Dasyatis bleekeri (Blyth) Journal of the American Oil Chemists Society 1998; 75(10):1373-1378
 - Dietary Guidelines for Americans. 2010. Washington, [22] DC: U.S. Government Printing Office.
 - Ould El Kebir, M.V., G. Barnathan, E.M. Gaydau, Y. Siau, [23] and J. Miralle's (2007). Fatty acid in liver, muscle, and gonad of three tropical rays including Non-Methylene-Interrupted Dienoic fatty acids. Lipids 42: 525-535
 - European Food Safety Authority (EFSA), 2010. [24] Scientific opinion on fish oil for human consumption. Food hygiene, including rancidity. The EFSA J.,8
 - [25] FAO. The Production of Fish Meal and Oil; Fishery Industries Division, Fish. Tech. Pap. 142, 1986; 63.
 - Farooqui, A. A. 2009. Beneficial effects of fish oil on [26] human brain. New York: Springer, pp .151-187.

- [27] Fei, C.Y., J. Salimon and M. Said. 2010. Optimisation of urea complexation by box behnken design. Sains Malaysiana, 39(5), 795-803.
- Frankel, E.N. 1998. Lipid oxidation. Chapter 10 foods. [28] The oily press, In: Frankel EN (ed). Dundee, UK, 187-225.
- Frankel, E.N. 1991. Recent advances in lipid oxidation. [29] Review. Journal of the Science of Food and Agriculture, 54, 495-511
- [30] Fritsche, K.L. and P.V. Johnston. 1988. Rapid autoxidation of fish oil in diets without added antioxidants. Journal of Nutrition, 118, 425-426.
- [31] Fry, J. P., & Love, D. C. 2013. Environmental Public Health and Recommendations for Fish Oil and SeafoodIntake. American Journal of Public Health, 103(11):e3-e4
- Gamez-Meza, N., J.A. Noriega-Rodri, L.A. Medina-[32] Juareza, J. Ortega-Garcaa, J. Monroy-Riverab, F.J. Toro-Vazquezc, H.S. Garcia and O. Angulo-Guerrerob. 2003. Concentration of eicosapentaenoic acid and docosahexaenoic acid from fish oil by hydrolysis and urea complexation. Food Research International, 36, 721-727.
- [33] Gardner, H.W. 1983. Effect of lipid hydroperoxides on food components. In xenobiotics in foods and feeds; Finley, J.W. Eds., American Chemical Society: Washington, DC, 63-84
- [34] *Garcia*-Martinez J, et al. (2004) The transcriptional response to transient cell wall damage on al JoA.I. Castello, and M.C. López-Sabater (2010).Long-chain in Saccharomyces cerevisiae and its regulation by the cell. Ciência Anim. Bras., 5 (3): 123-129
- Gray, J.I., E.A. Gomaa and D.J. Buckley. 1996. Oxidative arch an119: 54-61 [35] quality and shelf life of meats. Meat Science, 43, S111-0 [51] Moffat, C. F., & McGill, A. S. 1993. Variability of the S123
- [36] Gudmundur G. Haraldsson 1999 The application of 2456-647 lipase for preparing various lipids enriched with omega-3 fatty acids.Rit Fiskideildar 16.97-105
- Haagsma, N., C.M. Gent, J.B. Luten, R.W. Jong and E. [37] Doorn. 1982. Preparation of an omega-3 fatty acid concentrates from cod liver oil. Journal of American Oil Chemists' Society, 59, 117-118.
- [38] Harrison, K. 2007. EPA, Icosapentaenoic acid, Timnodonic acid, unsaturated fatty acid. http://www.3dchem.com/molecules.asp?ID=380.
- [39] Hayes, D.G., Y.C. Bengtsson, J.M. Van Alstine and F. Setterwall. 1998. Urea complexation for the rapid, ecologically responsible fractionation of fatty acids from seed oil. Journal of the American Oil Chemists' Society, 75, 1403-1409.
- [40] Helland I B, Smith L, Blomen B, Saarem K, Saugstad O D, Drevon C A.Effect of supplementing pregnant and lactating mothers with n-3 very long-chain fatty acids on children's IQ and body mass index at 7 years of age. Pediatrics. 2008;122: e472-9
- [41] Holub, D.J. and B.J. Holub. 2004. Omega-3 fatty acids from fish oils and cardiovascular disease. Molecular and cellular biochemistry, Kluwer academic publishers, Netherlands. 263, 217-225
- Huss, H.H. 1988. Fresh fish quality and quality changes. [42] Rome, Italy: FAO.

- Iverson, S.J., S.L.C. Lang and M.H. Cooper. 2001. [43] Comparison of the Bligh and Dyer and Folch methods for total lipid determination in a broad range of marine tissue. Lipids, 36, 11.
- Iqbal S. and M.I. Bhanger. 2005. Stabilization of [44] sunflower oil by garlic extract during accelerated storage. Food Chemistry.
- [45] Jayathilakan, K., G.K. Sharma, K. Radhakrishna and A.S. Bawa. 2007. Antioxidant potential of synthetic and natural antioxidants and its effect on warmedoverflavour in different species of meat. Food Chemistry, 105, 908-916
- Lands WEM. Fish and Human Health. Academic Press, [46] London, 1986, 1–186.
- [47] Lenihan-Geels, G.; Bishop, K.S.; Ferguson, L.R. 2013 Alternative Sources of Omega-3 Fats: Can We Find a Sustainable Substitute for Fish? Nutrients, 5, 1301-1315.
- Lehninger AL, Biochemistry, Edn 2, Worth Publishers.~ [48] 92

International Journal of Fisheries and Aquatic StudiesInc., NY, 1978, 279-301.

- [49] Liu, S., C. Zhang, P. Hong and H. Ji. 2006. Concentration of docosahexaenoic acid(DHA) and eicosapentaenoic acid (EPA) of tuna oil by urea complexation: • optimization of process parameters. Journal of Food Engineering, 73(3), 203-209
- global 🤍 [50] Mayneris-Perxachs, J., I. Bondia-Pons, L. Serra-Majem, n-3 fatty acids and classical cardiovascular disease risk factors among the Catalan population. Food Chem.,
 - composition of fish oils: significance for the diet.Proceedings of the Nutrition Society, 52(03), 441-456.
 - [52] Reuber, M.D. 1979. Carcinogenicity of chloroform. Environ. Health Perspect. 31, 171-182.
 - [53] Rizliya, V., & Mendis, E. 2014. Biological, Physical, and Chemical Properties of Fish Oil and Industrial Applications. In Seafood Processing By-Products, pp. 285-313.Springer, NewYork.
 - Navarro-Garcia, G., J.C. Ramirez-Suarez, E. Cota-[54] Quiñones, F. Márquez-Farías, and L. Bringas-Alvarado (2010).Storage stability of liver oil from two ray (Rhinopterabonasus and Aetobatusnarinari) species from the Gulf of Mexico. Food Chem. 119: 1578-1583.
 - Navarro-Garcia, G., R. Pacheco-Aguilar, B. Vallejo-[55] Cordova, J.C. Ramirez-Suarez, and A. Bolanos (2000).Lipid composition of the liver oil of shark species from the Caribbean and Gulf of California waters. J. Food Comp. and Analy. 13: 791-798.
 - [56] Navarro-Garcia, G., R. Pacheco-Aguilar, L. Bringas-Alvarado, and J. Ortega-Garci´a (2004). Characterization of the lipid composition and natural antioxidants in the liver oil of Dasyatis brevis and Gymnuramarmorata rays. Food Chem. 87: 89-96.
 - Norziah, M.H., J. Nuraini and K.Y. Lee. 2009. Studies on [57] the extraction and characterization of fish oil from wastes of seafood processing industry. Journal of the Science of Food and Agriculture, 2(4), 959-973

- [58] O'Brien, R.D. 2009. Fats and oils. Formulating and processing for applications, 3rd ed., CRC press, London, 213-300
- [59] Ould El Kebir, M.V., G. Barnathan, E.M. Gaydau, Y. Siau, and J. Miralle's (2007). Fatty acid in liver, muscle, and gonad of three tropical rays including Non-Methylene-Interrupted Dienoic fatty acids. Lipids 42: 525-535.
- [60] Ozogul, Y., A. Simşek, . Balikçi and M. Kenar. 2011. The effects of extraction methods on the contents of fatty acids, especially EPA and DHA marine lipids. in International Journal of Food Sciences and Nutrition
- Özyılmaz, A And Öksüz, A, 2015. Determination Of The [61] Biochemical Properties Of Liver Oil From Selected Cartilaginous Fish Living In The Northeastern Mediterranean. The Journal of Animal & Plant Sciences, 25(1): 2015, Page: 160-167 . ISSN: 1018-7081
- [62] Pak, C.S. 2005. Stability and quality of fish oil during typical domestic application. Final Project.Wonsan University of fisheries Kangwon Province, D.P.R. of Korea
- [63] Pasoz, M., J.M. Gallardo, J.L. Torres and I. Medina. 2005. Activity of grape poly phonesas inhibitors of the oxidation of fish lipids and frozen fish muscle. Food Chemistry, 92, 547-557
- [64] Pettinello, G., A. Bertucco, P. Pallado and A. Stassi. 2000. Production of EPA enriched mixtures by supercritical fluid chromatography: from the laboratory scale to the pilot plant. Journal of Supercritical Fluids, 19, 51-60
- [65] Ramakrishnan U, Stein AD, Parra-Cabrera S, Wang M, Imhoff-Kunsch B, Juarez-Marquez S, Rivera J, Martorell R. Effects of ar [79] Sun Y, et al. (2005) Interaction of Sla2p's ANTH domain docosahexaenoic acid supplementation during opmen pregnancy on gestational Vage and size at birth: randomized, double-blind, placebo-controlled [80] trial in Mexico. Food Nutr Bull. 2010;31:S108–16
- Ratanyake, W.M.N., B. Olsson, D. Matthews and R.G. [66] Ackman. 1988. Preparation of omega-3 PUFA concentrates from fish oil via urea complexation. Fat ScienceTechnology, 90, 386-391.
- Raza, S.A., A. Rashid, J. William, S. Najaf and M. Arshad. [67] 2009. Effect of synthetic antioxidant on shelf life of locally manufactured butter known as Makhan in Pakistan. Biharean Biologist, 3(2), 161-162
- Rubio-Rodríguez, N., S.M. de Diego, S. Beltrán, I. Jaime, [68] M.T. Sanz and J. Rovira. 2008. Supercritical fluid extraction of the omega-3 rich oil contained in hake (Merluccius capensis-Merluccius paradoxus) byproducts: study of the influence of process parameters on the extraction yield and oil quality. Journal of Supercritical Fluids, 47, 215-226
- [69] Saify, Z.S., S. Akhtar, K.M. Khan, S. Perveen, S.A.M. Ayattollahi, S. Hassan, M. Arif, S.M. Haider, F. Ahmad, S. Siddigui and M.Z. Khan. 2003. A Study on the fatty acid composition of fish liver oil from two marine fish Eusphyra blochii and Carcharhinus bleekeri. Turkish Journal of Chemistry, 27, 251-258
- [70] Sang, W. and Z.T. Jin. 2004. Lipid oxidation of fish liver affected light, oil by as

antioxidants and temperature. Journal of Food Processing and Preservation, 28(1), 1-10.

- S.P.J.N. and F. [71] Senanayake, Shahidi. 2000. Concentration of DHA from algal oil via urea complexation. Journal of Food Lipids, 7, 51-61.
- Simopoulos, A. P. (1991). Omega-3-fatty-acids in health [72] and disease and in growth and development. American Journal of Clinical Nutrition, 54(3), 438-463.
- [73] Simopoulos, A.P (2006). Evolutionary aspects of diet, theomega-6/omega-3ratio and genetic variation: nutritional implications for chronic diseases, Biomedicine and Pharmacotherapy, vol. 60, no. 9, pp. 502-507.
- Simic, M.G., K.A. Taylor, J.F. Ward and C. Von Sonntag. [74] 1988. Oxygen radicals in biology and medicine, Plenum Press, New York, NY.
- Smedes, F. and T.K. Askland. 1999. Revisiting the [75] Development of the Bligh and Dyertotal lipid determination method, Marine Pollution Bulletin, 38, 193-201.
- [76] Stansby ME. Fish oils, their Chemistry, Technology, Stability, Nutritional properties and Uses, Avi Publishing Company. Inc 1968, 3-9
- Strlič, M., I.K. Cigić, I. Rabin, J. Kolar, B. Pihlar and M. E[77] Cassar. 2009. Autoxidation of lipids in parchment, polymer degradation stability, 94, 886-890
 - Strocchi, A. and G. Bonaga. 1975. Correlation between [78] urea inclusion compounds and comformational structure of unsaturated C-18 fatty acid methyl esters. Chemical Physical Lipids, 15, 87-94.
 - with PtdIns(4,5)P2 is important for actin-dependent endocytic internalization. Mol Biol Cell 16(2):717 30
- Tor, C. and H. Yi. 2001. Polyunsaturated fatty acid concentrates from borage and linseed oil fatty acid. Journal of America Oil Chemists' Society, 78(5), 485-.... 488.
 - [81] Uauv R, Valenzuela A 2000 .Marine oils: the health benefits of n-3 fatty acids. Nutrition.;16(7-8):680-4.
 - Undeland, I. 1998. Lipid oxidation in fillets of herring [82] (Clupea harengus) during processing and storage. Ph.D. Thesis, Chalmers University of Technology, Goteborg, Sweden
 - [83] Wang, C., Harris, W.S., Chung, M., Lichtenstein, A.H., Balk, E.M., Kupelnick, B., Jordan, H.S., Lau, J. 2006. n-3 Fatty acids from fish or fish-oil supplements, but not alpha-linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review. American Journal of Clinical Nutrition 84 (1): 5–17.
 - Wanasundara, U.N. and F. Shahidi. 1998. Lipase-[84] assisted concentration of omega-3 polyunsaturated fatty acids in acylglycerol form from marine oils. JournalAmerican Oil Chemists' Society, 75, 945-95.
 - Wanasundara, U.N. and F. Shahidi. 1999. Concentration [85] of omega 3-polyunsaturated fatty acids of seal blubber oil by urea complexation: optimization of reaction conditions. Food Chemistry, 65, 41-49.
 - World Health Organization.1982. Expert Committee on [86] the Prevention of Coronary Heart Disease, WHO Tech. Rep. Ser. 678, Geneva, Switzerland.