Virtual Reality as a Promising Tool for Autism Intervention

Sinitha. K. M (MASLP)¹, Stephy Jacob (MASLP)², Dr. Maria Grace Treasa PhD (SLP)³

¹Audiologist and Speech Pathologist: Department of Audiology & Speech Language Pathology National Institute of Physical Medicine & Rehabilitation (NIPMR), Irinjalakuda, Kerala, India ²Audiologist and Speech Pathologist: Private Practitioner, Thrissur, Kerala, India ³Paediatric Speech - language Pathologist: Aster KIND (Kids Integrated Neuro Developmental Center) Aster Medcity, Kochi, Kerala, India

ABSTRACT

This paper explores how Virtual Reality (VR) systems have been used as a rehabilitation tool for disabled population. Reviews were done in applications of virtual reality in patients with neurological disorders, visual impairment, psychiatric problems, children with physical disability and neurodevelopment disorders. This article mainly focused on the current status and use of virtual reality for children with autism. Literature was reviewed and the important findings are discussed in this paper. The virtual reality systems & designs, interventions method, treatment intensity and its effectiveness in target population were analyzed. The following skills emotion recognition, contextual processing, social attribution & executive function of analogical reasoning, navigation performance, safety skills, social interaction, motor & cognitive skills, conversational understanding were found to be improved in children with autism spectrum disorder by using VR. The paper also detailed the studies done in India using virtual reality in the disability field, mainly in autistic population.

KEYWORDS: Virtual Reality, Autism Spectrum Disorder, Rehabilitation tool, Disability

of Trend in Scientific Research and Development

INTRODUCTION

Virtual reality (VR) is a promising technology which has applications in wide range of aspects. A systematic literature review was done in this paper to determine the uses of VR in the field of rehabilitation especially in children with autism. This work also investigated the status of VR technology in the rehabilitation of individuals with autism in Indian scenario.

Virtual reality has been defined as an "interactive, immersive experience generated by a computer" (Pimentel & Teixeira, 1993).Human-computer interactivity is attained through multiple sensory channels which allow people to explore Virtual Environments (VE) through sight, sound, touch and sometimes even smell (Wang & Reid, 2011).In terms of human experience VR is a mediated environment which creates the sensation in a user of being present in a physical surrounding (Seipel, 1993).

Virtual reality is about generating "acceptable "reproductions of real things or environments for training, entertainment or design purposes. It uses computers to generate 3D environments in which one can navigate and interact. Navigation implies the capability to move around and explore the features of a 3D scene, such as walking through a forest. And interaction means the capacity to select and manipulate things in the scene, for example, grabbing and examining a flower found in the forest. In other words, VR fully immerses the user in a three-dimensional, computer-generated world and depending on the system *How to cite this paper:* Sinitha. K. M | Stephy Jacob | Dr. Maria Grace Treasa "Virtual Reality as a Promising Tool for

Autism Intervention" Published in International Journal of Trend in Scientific Research and Development (ijtsrd), ISSN: 2456-6470, Volume-5 | Issue-4,



June 2021, pp.1274-1280, URL: www.ijtsrd.com/papers/ijtsrd42523.pdf

Copyright © 2021 by author (s) and International Journal of Trend in Scientific Research and Development Journal. This is an Open Access article distributed

under the terms of the Creative Commons Attribution License (CC



License (CC BY 4.0) (http://creativecommons.org/licenses/by/4.0)

used; the amount of immersion in the virtual world can be any place from barely being capable to affect virtual objects to feeling every part of the virtual world. The target of anyone involved in the field of VR is to create an environment that is indistinguishable from the real world. It requires real-time graphics, a stereoscopic display, used to produce the illusion of 3D and a tracking system to obtain head and hands motion. Head-Mounted Displays and Stereoscopic glasses are commonly used technologies (Gutierrez, F, & Thalmann, 2008).

There are different types of VR systems. A major difference between these VR systems is the mode with which they interface to the user. Some of the common modes used in VR systems includes; Window on World Systems/Desktop VR, Video Mapping, Immersive Systems, Telepresence and Mixed Reality (Isdale, 1993).

Clinical Applications of VR

Virtual reality technology has broad range of applications in the field of rehabilitation. VR has proven its applications in patients with neurological disorders, visual impairment, psychiatric problems and children with physical disability and neurodevelopmental disorders.

Benefits of VR for children with disabilities:

In recent years, VR tools have originated as an assessment and treatment methods in rehabilitation of disabled population. The properties of VR, especially its malleability, make it a flexible medium for the creation of customized applications to satisfy the requirements of disabled children. Depending on the treatment goal of program the VE characteristics can be modified to incorporate or exclude certain categories of stimuli. This adaptability helps to encourage an optimal interaction for these children. VR can be considered as an assistive technology, due to its countless potential to reduce the effects of disability in these populations. VR technology also offers children with disabilities, larger access to many experiences and can enhance various skills (Mccomas, Pivik, & Laflamme, 1998).

A variety of unique attributes of VR technology are the rationale for using it in rehabilitation field. These distinctive attributes include the opportunity for experimental and active learning which inspires and motivates the patient. Additionally there is the capability to objectively measure patient's behavior in challenging environments. The stimulus delivery and its measurement can be controlled experimentally. VEs provide the chance for repeated learning in different trials and provide the capacity to increase the task complexity gradually while fading the help offered by therapist. The stimulus delivery in VE is automated in nature which helps the therapist to provide physical support according to patients need without detracting task complexity (Weiss, Kizony, Feintuch, & Katz, 2006).

Applications in neurorehabilitation:

Virtual Reality applications in the rehabilitation of neurologically impaired population are expanding at a fast pace and a huge variety of platforms and programs have been developed. Its application is evident in patients with stroke and traumatic brain injury(Weiss, Kizony, Feintuch, & Katz, 2006).The results of the VR technology study for physical rehabilitation in patients with chronic frozen shoulder and post-traumatic brain injury suggested that effective VR-based rehabilitation could be used as a supplement or alternative to conventional therapy (McComas, et al., 2003).

In stroke patients, VR can be used as a more comprehensive and controlled evaluation tool of prospective memory than other standard memory tests (Brooks, Rose, Potter, Attree, Jayawardena, & Morling, 2002). For individuals with unilateral spatial neglect due to right hemisphere stroke,non- immersive interactive VEs are an effective medium for teaching the safe road crossing task (Naveh, Katz, & Weiss, 2000).

Applications in psychiatry:

Patients with psychiatric illness were also benefitted by using VR technology. In schizophrenia patients, social skills training using VR role-playing were compared with traditional role –playing. It was found that VR may be a helpful supplement to traditional social skill training because of the improvement noticed in terms of conversational skills and assertiveness (Kyung-MinPark, et al., 2011).

A study result using a application of VR integrated program showed improvements in schizophrenia patient's social dysfunction behaviors like negative symptoms, anxiety, discomfort and avoidance and thus it can be used as an adjunct technique for their social skill intervention(Calafell, Maldonado, & Sabate, 2014).

Applications in visual impairment:

Applications of VR in visually impaired people are also emerging. An application called Virtual Reality Simulator which is capable of making an auditory representation of VE was developed for visually impaired people. It helps in rendering the virtual world completely through hearing (Torres-Gil, Gonzalez, & Gonzalez Mora, 2010).

Applications in neurodevelopmental disorders:

Aguirre et al (2019) carried out a project by creating a VR game, which places intellectually disabled students to become very confident when they are faced with a stressful scenario in their everyday lives. The VR environment make a safe space in which they can develop their social abilities and increase their situational awareness, hence they can train themselves to be more relaxed when they face stressful situations in the true world (Aguirre, et al., 2019).

Studies were conducted using technology based games such as Wheelchair Net and Social MatchUP. The contribution of Wheelchair Net, a technology based training program in the education of physical disability children was studied by giving a chance to practice driving in virtual motorized wheelchairs in a computer-generated world safely. The study results showed achievement in driving skills of participants as indexed in true reality (Inman, Loge, Cram, & Peterson, 2011). Social MatchUP, is a multiplayer VR game for children with neurodevelopmental disorders which enhances their social and communication skill, and it can also be used as a learning tool. It is a simple concentrationlike game, run on smartphones (Loiacono, Trabucchi, Messina, Matarazzo, Garzotto, & Beccaluva, 2018).Vona et.al (2020) did a case study by means of the Social Matchup application for people with neurodevelopmental disorders to develop their speech-based conversational capability. The participants were 24 persons in 2 groups, one group of participants playing a game in Social MatchUP and the other group playing a game that is similar as in the application but in the real world. Their results point out that the game experience in Social MatchUP was usable and enjoyable, and boosted top conversational abilities with respect to its corresponding item in the physical setting (Vona, Silleresi, Beccaluva, & Garzotto, 2020).

There is paucity of **Indian studies** on use of VR for rehabilitation. Reviewed studies are summarized as follows,

In a study by Doctor et.al (2019) was found that VR training to be therapeutic in improving psychosocial factors of spinal cord injury patients. Their reduction in stress, anxiety and increased calmness were reported in post intervention (Doctor, Gadgerao, Udpikar, Shyam, & Sancheti, 2019).A systematic review was done by Ravi et.al (2017) to provide updated evidence-based guidelines for VR rehabilitation of patients with cerebral palsy in their sensory and functional motor skills. They concluded that VR rehabilitation is a promising intervention for improving balance and motor skills of patients with cerebral palsy (Ravi, Kumar, & Singhi, 2017). A study was done by Sharan et.al (2012) to find out the effectiveness of VR based training in the rehabilitation of children with cerebral palsy. There were 14 participants in the study group and 15 subjects in the control group. Outcome measures used in the study were Manual Ability Classification System, Pediatric Balance Scale, child satisfaction, level of participation, motivation and cooperation. While comparing with the control group, the results of study group were reported to be improved significantly in all the outcome measures except in Manual Ability Classification System (Sharan, Ajeesh, Rameshkumar, Mathankumar, Paulina, & Manjula, 2012).

Muneer et.al (2015) explored the use of VR-based games to improve the motor, communication, cognitive and social/emotional skills in children with developmental

disabilities. Their result reveals significant progress in motor, cognitive and social/emotional skills (Muneer, Saxena, & Karanth, 2015).

Based on the reviews done above, areas for which VR has been successfully used in the rehabilitation field is summarized in the following Table.

SL NO	AUTHORS	YEAR	AREAS FOR WHICH VR HAS BEEN USED	SUBJECT / POPULATION STUDIED					
1	Brooks, Rose, Potter, Attree, Jayawardena, & Morling.	2002	Prospective memory assessment	Stroke					
2	Naveh, Katz, & Weiss	2000	Safe road crossing task	Unilateral spatial neglect					
3	Kyung-MinPark, et al	2011	Social skills	Schizophrenia					
4	Calafell, Maldonado, & Sabate	2014	Social skills	Schizophrenia					
5	Torres-Gil, Gonzalez, & Gonzalez Mora	2010	Environment recognition, auditory rendering	Visual impairment					
6	Inman, Loge, Cram, & Peterson,	2011	Driving skills	Physical disability					
7	Aguirre, et al	2019	Social skills and situational awareness	Intellectual disability					
8	Loiacono, Trabucchi, Messina, Matarazzo, Garzotto, & Beccaluva	2018	Social and communication skill	Neurodevelopmental Disorders					
9	Vona, Silleresi, Beccaluva, & Garzotto	2020	Conversational ability	Neurodevelopmental Disorders					
10	Doctor, Gadgerao, Udpikar, Shyam, & Sancheti	2019	Psychosocial factors	Spinal cord injury					
11	Ravi, Kumar, & Singhi 🥢 🌽 🥂	2017	Balance and motor skills	Cerebral palsy					
12	Deepak, P S, R, M, Jospin. 🥖 🛸 🔹	2012	Balance skills	Cerebral palsy					
13	Muneer, Saxena, & Karanth	2015	Motor, cognitive and social/emotional skills	Developmental disabilities					

Table 1: Successful use of VR in the rehabilitation field

Application of VR in Autism

This article has a robust literature review regarding the present status and use of VR for autism spectrum disorder. Autistic individuals have impairments in social and communication skills, lack of imagination or abstract thought and the presence of stereotyped, repetitive behaviors. Considering the balance between flexibility and control of the treatment program in autistics, VR systems play an excellent role in treatment outcome. Several skills from motor, cognitive, social and emotional domains were investigated in these studies. Autistic individuals improved their learning and transferred to new skills in considerably less time through VR systems.

Table 2

A detailed literature review was done on the uses of VR in autism and is summarized in the table below.

Source	Participant characteristics	Description of treatment	Description of results				
Yang et al(2017)	Sample size:	Design: Within subjects	VR-SCT significantly				
	17young adults	VR System: Windows computer	improved emotion				
	Gender: 2 females,	Webcam and a headphone with built-in	recognition in terms of the				
	15 males	microphone.	ACS-SP scaled scores and in				
	Mean age: 22.50	Intervention: Virtual Reality Social	theory of mind from				
	years	Cognitive Training (VR-SCT).	pretreatment to post				
	Diagnosis- : High	Treatment intensity: 2x 1 hour session	treatment.				
	functioning Autism	per week for 5-weeks					
Wang et al(2010)	Sample size: 4	Design: single subject design with	Improvements in contextual				
	children with	nonconcurrent multiple baselines	processing ability from				
	autism	across subjects	baseline to treatment for each				
	Gender: 3 boys, 1	VR System: laptop computer with	child, with average increases				
	girl	tracking web cam	from 15% to 46%. All				
		Intervention: Virtual Reality-Cognitive	children maintained a high				
	Age range : 5-10	Rehabilitation (VR-CR).	level of performance at the				
	years	Treatment intensity: baseline phase: 4	two-week follow-up				
	Diagnosis: Autism	to 6 weeks	assessment.				
	spectrum disorder	Treatment phase: 3 discrete sessions x					
		10 minutes					

International Journal of Trend in Scientific Research and Development (IJTSRD) @ www.ijtsrd.com eISSN: 2456-6470

Sait et al (2019)	Sample size: 9 children	Design: within subject study VR system: AutiVE, VR glasses,	VR system helps the autistic child's to enrich the ability to
	Gender: not	interfaces from website and unity	adjust in a new environment.
	specified	scenes	8 children were satisfied and
	Age range : 4-12	Intervention: VRinto the educational	accept the VR headset. 5 of
	years old	system(school scene and classroom	them remembered the scene
	Diagnosis: Autism	scene)based on applied behavior	presented. Few words were
Didahbani at al(2016)	Sample size: 20	Decign: within subjects	Significant improvements
Didenbani et al(2010)	children	VR System: Second Life™ version 2.1	were noticed on emotion
	Gender:	three-dimensional virtual world	recognition social attribution
	26 males.4 females	software (Linden Lab. 2003). Microsoft	attention and executive
	Age range : 7-16	Windows XP or newer, graphics cards	function of analogical
	vears	of ATI Radeon 8500 or better and 1.5	reasoning.
	Diagnosis: Autism	GHz 86 CPU using a 24-inch monitor	5
	spectrum disorder	with a resolution of 1920 x 1200.	
		Intervention: Virtual Reality Social	
		Cognition Training Treatment intensity:	
		10x 1 hour sessions across 5 weeks	
	Sample size: 7	Design: within subjects	Over sessions, 6 out of 7
Saiano et al(2015)	adults	VR System: video projector, displaying a	significantly improved their
	Gender: allmale	VR environment on a 2 m × 2 m screen,	navigation performance, but
	Age range : 19-44	Markerless motion capture device	did not reduce the errors
	years	(MICrosoft Kinect)	Both parents and carogivers
	spectrum disorder	Toolkit (EAAST)	stated a significant progress
	spectrumusoruer	Intervention: integrated approach	in the subject's street crossing
	AN	hased on VE and natural interfaces (ahility
	AN	Street crossing)	Interaction with VEs effective
	A S.	Treatment intensity: 10x45 minutes	in facilitating the acquisition
	2 2 .	sessions(1session/week)	of safety skills in autism
Self et al(2007)	Sample size: 8	Design: between groups 👔 🔹 💋	Both groups improved in
	children 💋 🧧 🎍	(randomly assigned to 2 groups)	their learning and transfer of
	Gender : 6 boys 2	VR System: laptop computer with a	safety skills. The VR group,
	girls Age range: 6	CPU, 3DSMax6.0 software, a separate	however, learned these skills
	to 12 years	19-in Ultra Sharp Dell flat panel	in considerably less time.
	Diagnosis : Autism	monitor ,a Scent Palate	
	spectrum disorder	Intervention: comparing benefits of	
	Nr.	using VR to benefits of an	
	Y V	safety skills Treatment intensity: 2x30	
		minutes for 5 week for each phase	
Yuan et al(2018)	Sample size: 72	Design: within subjects	Children from training group
	children	VR System: four-side Cave Automatic	scored higher on emotion
	Gender: 64 boys, 8	Virtual Environment (CAVE),goggles	expression, regulation, social
	girls	Intervention: VR enabled training	interaction and adaptation
	Mean age: 106.3	program. Six learning scenarios were	after the training than before
	months	designed, including a relaxation	the training. There was a
		scenario, four training scenarios and	statistically significant
		one consolidation scenario.	interaction between group
		Treatment intensity: The one-hour	and time on affective
		training session would end with a	expressions and on social
Karanth at al(2015)	Sample size: F	Design: within subjects	Pro and post intervention
Karanth et al(2015)	sample size: 5	VP System: Yboy260(dovoloped by	scores revealed that the
	Gender: 4 hove 1	Microsoft consists of a console with a	children made significant
	girl.	240 GB hard drive and a controller) and	progress, not only in certain
	Age range: 4-8	Kinect (motion-sensing device created	motor skills but also in skills
	years	by Microsoft)	from the cognitive and
	, Diagnosis : 4 bovs	Intervention: VR Games	social/emotional domains.
	with Autism	Treatment intensity: 4 and 6 sessions	None of the children
	Spectrum Disorder	(20-30 minutes) over a span of one	regressed in any of the skills
	and 1 girl with	month.	monitored from the different
	Learning Disability		domains.
Rosenfield et al(2019)	Sample size:	Design: within subjects	Through the repetition and

International Journal of Trend in Scientific Research and Development (IJTSRD) @ www.ijtsrd.com eISSN: 2456-6470

2children Gender :	VR System: Oculus Rift SDK,Oculus Rift	analysis of these virtual
1boy, 1 girl .	VR headset,VE comprises five primary	interactions, users improved
Mean age: 6.5	software modules: Staging script, vision	social and conversational
years	processing script, voice processing	understanding.
Diagnosis: Autism	script, data archive script, and character	The study was a positive
Spectrum Disorder	animations.	experience for the users, and
	Intervention: VR-based assistive	minimal training was
	technology which integrates gaze	required to use the VR
	tracking and voice processing	headset. The users quickly
	Treatment intensity: 15 nonconsecutive	discovered the immersive
	minutes	nature and malleability of the
		system, while also interacting
		socially in the physical world.

Table 2: Use of VR in autism

VR Studies in India on Autistic Population

Review of literature reveals few Indian studies on VR in ASD.

A virtual environment based therapy was proposed by Manju et.al, (2018) to enhance the social skills, emotional skills and attention of the children with autism. The virtual environment included 3 levels, first level focused on increasing attention, second level focused on improving social interaction and third level focused on enhancing decision making. The proposed therapy produced positive results on repetition and it also noticed at what stage these children became panic, frustrated and enthusiastic (Manju, Padmavathi, & Tamilselvi, 2018). VR based joint attention task platform was developed by Jyoti et.al (2020) which was augmented with hierarchical prompt protocol such as using eye cue, head turn, finger pointing, and sparkling cues. The participants included in the study were 20 pairs of agematched autism spectrum disorder and typically developing children and the study results indicates the potential of the system to identify their deficits in joint attention skills. The study group showed difficulty in following eye cue, the ability to follow finger pointing was the best, and some could pick up merely the sparkling cue. All typically developing participants were able to follow eye cue without requiring other cues (Jyoti & Lahiri, 2020).

A physiology-sensitive VR-based social communication system (which is equipped with anxiety-sensitive feature) was developed by Kuriakose et.al (2017) to improve autism children's performance in a social task. The equipment can objectively identify and quantify one's anxiety level by monitoring physiological biomarkers thus helps to offer tasks of varying challenge levels in an individualized way thereby helping to encourage improved social communication-related performance. This can help to recognize elements of social interaction which is anxietyprovoking for them. Thus in future this system may help to build up a comprehensive social ability learning platform that can improve quality of life of persons with autism (Kuriakose & Lahiri, 2017). The effect of using collaborative VEs during intervention programs for children with autism was studied by Yogeswara et.al (2013). There were 10 subjects in both the experimental and control group. The two tasks given in the study were emotion recognition and theory of mind. In emotional recognition task the subjects were asked to respond only for happy faces when both the neutral and happy faces are presenting. And in the theory of mind task participants were asked to make two judgements; reality judgment and a think judgment. Studies were done during the pre and post intervention stage and compared responses with and without collaborative VEs. Results imply that the presentation of emotional expressiveness as part of

VE helped autistic children to communicate without fear and thus recognize others emotions (Yogeswara, Kumar, Santosh, & Anand, 2013).

A study was proposedby Shree et.al (2020) which contributes the children affected with autism to develop managing their safety awareness on fire by recognizing the environment that is about to cause a fire disaster and help them to make exit from the situation. They generated a model environment to create awareness towards the safety on fire by finishing the tasks given through stages and to observe them to examine their task performances and formulate the participants to energetically immerse into virtual experiences. Thus this system helped children with autism to learn about fire cautions, how to react and get away from a fire accident (Shree & Selvarani, 2020).

The literature review of Indian studies on persons with autism indicates that VR can be used in various domains such as to identify deficits in joint attention, intervene on social communication, safety and following social rules.

Conclusion

VR technology has abilities and its benefits in the rehabilitation field mainly in autism are evident from the reviews done in this paper. From the review of Indian studies in autism, it is clear that most of the studies were happened in IIT rather than in a rehabilitation setup. However those study results also revealed positive outcomes in children with autism. Thus in future more studies should occur in India in the field of autism rehabilitation using VR. VR has a huge potential to become a promising tool for autism intervention.

BIBLIOGRAPHY

- [1] Aguirre, C., Angulo, M., Barkley, M., Evans, G., Fitzsimmons, C., Vu, M., et al. (2019). Virtual Reality Program To Help Intellectually Disabled Students Improve Social Skills And Situational Awareness. *Assistive Technology-RESNA ANNUAL CONFERENCE* (pp. 231-250). Arlington: RESNA.
- [2] Brooks, M. B., Rose, F. D., Potter, E. A., Attree, S., Jayawardena, S., & Morling, A. (2002). Assessing stroke patient's ability to remember to perform actions in the future using virtual reality. *proceedings of the 4th international conference on disability*, *Virtual reality and associated technology*, (pp. 239-245).
- [3] Calafell, M., Maldonado, J., & Sabate, J. (2014). A virtual reality-integrated program for improving social skills in patients with schizophrenia: A pilot

[22]

study. *Journal of Behavior Therapy and Experimental Psychiatry*, 45 (1), 81-89.

- [4] Doctor, K., Gadgerao, S., Udpikar, R., Shyam, A. K., & Sancheti, P. K. (2019). Effect of virtual reality training on psychosocial factors in spinal cord injury patients. *International Journal of Health Sciences & Research*, 9 (8), 433-438.
- [5] Gutierrez, M., F, V., & Thalmann, D. (2008). *Stepping into Virtual Reality.* Springer Science & Business Media.
- [6] Inman, D. P., Loge, K., Cram, A., & Peterson, M. (2011). Learning to Drive a Wheelchair in Virtual Reality. *Journal of special education technology*, 26 (3), 21-34.
- [7] Isdale, J. (1993). What is Virtual Reality? Retrieved october 13, 2020, from vr.html: http: //www.columbia/~rk35/vr/vr.html
- [8] Jyoti, V., & Lahiri, U. (2020). Virtual Reality Based Joint Attention Task Platform for Children With Autism. *IEEE Transactions on Learning Technologies*, *13* (1).
- [9] Kuriakose, S., & Lahiri, U. (2017). Design of a Physiology-Sensitive VR-Based Social Communication Platform for Children With Autism. *IEEE Trans Neural Syst Rehabil Eng*, 25 (8), 1180-1191.
- [10] Kyung-MinPark, JeonghunKu, HeeChoi, S., JeongJang, H., YeonPark, J., Kim, S. I., et al. (2011). A virtual reality application in role-plays of social skills training for schizophrenia: A randomized, controlled trial. *Journal of Behavior Therapy and Experimental Psychiatry*, 45 (1), 81-89.
- [11] Loiacono, T., Trabucchi, M., Messina, N., Matarazzo, V., Garzotto, F., & Beccaluva, E. A. (2018). Social MatchUP
 -: a Memory-like Virtual Reality Game for the Enhancement of Social Skills in Children with Neurodevelopmental Disorders. *Extended Abstracts of the 2018 CHI Conference* (pp. 2-7). canada: [24] ResearcGate.
- [12] Manju, T., Padmavathi, S., & Tamilselvi, D. (2018). A Rehabilitation Therapy for Autism Spectrum Disorder Using Virtual Reality. *International Conference on Intelligent Information Technologies* (pp. 328-336). Singapore: Springer.
- [13] Mario Saiano, Laura pellegriano, Maura Casadio, Susanna Summa, Eleonora Garbarino, Valentina Rossi,Daniela Dall Agapa, Vittorio Sanguineti. (2015). Natural interfaces and virtual environments for the acquisition of street crossing and path following skills in adults with ASD: a feasibility study. *Journal of neuro engineering and rehabilitation*, 12-17.
- [14] Mccomas, J., Pivik, J., & Laflamme, M. (1998). Current Uses of Virtual Reality for Children with Disabilies. *Virtual Environments in Clinical Psychology and Neuroscience*, 1-10.
- [15] McComas, J., Thornton, M., Marshall, S., Finestone, H., McCormick, A., Mayhew, A., et al. (2003). Experimental studies of virtual reality-delivered compared to conventional exercise programs for rehabilitation. *cyberpsycholBehav*, 6, 245-249.

- [16] Muneer, R., Saxena, T., & Karanth, P. (2015). Virtual Reality Games as an Intervention for Children: A Pilot Study. *Disability, CBR & Inclusive Development, 26* (3), 77-96.
- [17] Nalalia Stewart Rosenfield, Kathleen Lamkin, Jennifer Re, Kendra Day, Lou Anne boyd, Erik Linstead. (2019).
 A VR system for practicing conversation skills for children with autism. *Multimodel technologies and interaction, 3, 28.*
- [18] Naveh, Y., Katz, N., & Weiss, P. L. (2000). The effect of interactive virtual environment training on independent safe street crossing of right CVA patients with unilateral spatial neglect. *3rd international conference on disability,virtual reality and associated technology*, (pp. 243-248). italy.
- [19] Nyaz Didehbani, Tandra Allen, Michelle Candalaft, Daniel Krawczyk, Sandra Chapman . (2016). social cognition training for children with high functioning autism. *computers in human behaviour*, *62*, 703-711.
- [20] Pimentel, K., & Teixeira, K. (1993). *Virtual Reality through the New Looking Glass.* Windcrest, NY.
- [21] Ravi, D., Kumar, N., & Singhi, P. (2017). Effectiveness of virtual reality rehabilitation for children and adolescents with cerebral palsy: an updated evidence-based systematic review. *Physiotherapy*, *103* (3), 245-258.
 - Sait, M., Alattas, A., omar, A., Almalki, S., Sharf, S., & Alsaggaf, E. (2019). Employing Virtual Reality techniques in environment adaptation for autistic children. *Procedia computer science*, *163*, 338-344.
- [23] Seipel, S. (1993). An Introduction into Virtual Reality Environments. Retrieved October 13, 2020, from Microsoft PowerPoint-F1_070928_Intro_IGS: https: //www.it.uu.se/edu/course/homepage/igs/ht06/lec /F1 igs intro vr.pdf
- [24] Sharan, D., Ajeesh, P. S., Rameshkumar, R., Mathankumar, M., Paulina, R. J., & Manjula, M. (2012).
 13. Virtual reality based therapy for post operative rehabilitation of children with cerebral palsy. *Work*, 41, 3612-3615.
- [25] Shree, T. N., & Selvarani, A. G. (2020). Virtual Reality based System for Training and Monitoring Fire Safety Awareness for Children with Autism Spectrum Disorder. 5th International Conference on Devices, Circuits and Systems. IEEE.
- [26] Sze Ngar Vanessa Yuan, Horace Ho Shinglp. (2018). Using virtual reality, train emotional and social skills in childern with AST. *London journal of primary care*, *10*, 110-112.
- [27] Torres-Gil, M. A., Gonzalez, O. C., & Gonzalez Mora, J. L.
 (2010). Applications of virtual reality for visually impaired people. WSEAS transactions on computers, 9
 (2), 184.
- [28] Trisha Self, Rosalind R Scudder, Gamal Weheba, Daiquirie Crumrine. (2007). Avirtual approach to teaching safety skills to children with AST. *Top lang disorders, 27,* 242-253.
- [29] Virtual Reality Program To Help Intellectually Disabled Students Improve Social Skills And

Situational Awareness., (pp. C. . Angulo 1, M. Barkley 1, G. Evans 1, C. Fitzsimmons 1, M. Vu 1, N. Peixoto 1).

- [30] Vona, F., Silleresi, S., Beccaluva, E., & Garzotto, F. (2020). Social MatchUP: Collaborative Games in Wearable Virtual Reality for Persons with Neurodevelopmental Disorders. *Joint International Conference on Serious Games* (pp. 49-65). switzerland: Springer International Publishing.
- [31] Wang, M., & Reid, D. (2011). Virtual Reality in Pediatric Neurorehabilitation: Attention Deficit Hyperactivity Disorder, Autism and Cerebral Palsy. *Neuroepidemiology*, *36*, 2-18.
- [32] Wang, M., & Reid, D. (2013). Using the virtual reality cognitive rehab approach to improve contextual

processing in children with autism. *The scientific world journal*, 9 pages.

- [33] Weiss, P. L., Kizony, R., Feintuch, U., & Katz, N. (2006). Virtual reality in neurorehabilitation. *Textbook of Neural Repair and Neurorehabilitation*, 187-192.
- [34] Yang, Y. J., Allen, T., Abdullahi, S. M., Pelphery, K. A., Volkmar, F. R., & Chapman, S. B. (2017). Brain response to biological motion predict treatment outcome in young adults with autism receiving virtual reality social cognition training: preliminary findings. *Behaviour Research and Therapy*, 93, 55-66.
- [35] Yogeswara, R. M., Kumar, S., Santosh, J., & Anand, S. (2013). Virtual Technologies in Intervention Programs for Autistic Children. *International Journal* of Emerging Technologies, 4 (1), 39-43.

